



K24P 0864

Reg. No. :

Name :

**Second Semester M.Sc. Degree (C.B.S.S. – Supple. (One Time Mercy
Chance)/Imp.) Examination, April 2024
(2017 to 2022 Admissions)**

MATHEMATICS

MAT 2C 09 : Foundations of Complex Analysis

Time : 3 Hours

Max. Marks : 80

PART – A

Attempt **any four** questions from this part. **Each** question carries **4** marks :

1. Given that γ and σ are closed rectifiable curves having the same initial points. Prove that $n(\gamma + \sigma, a) = n(\gamma, a) + n(\sigma, a)$ for every $a \notin \{\gamma\} \cup \{\sigma\}$.
2. Let f be analytic on $B(0, 1)$ and suppose $|f(z)| \leq 1$ for $|z| < 1$. Show that $|f'(0)| \leq 1$.
3. Does the function $f(z) = z^2 \sin\left(\frac{1}{z^2}\right)$ has an essential singularity at $z = 0$? Justify your answer.
4. Using residue Theorem, prove that $\int_0^\infty \frac{1}{1+x^2} dx = \frac{\pi}{2}$.
5. Define the set $C(G, \Omega)$ and show that it is non-empty.
6. State the Weierstrass Factorization theorem.

PART – B

Answer **any four** questions from this part without omitting any Unit. **Each** question carries **16** marks :

Unit – I

7. a) Prove the following : If G is simply connected and $f : G \rightarrow \mathbb{C}$ is analytic in G then f has a primitive in G .
b) State and prove The Open Mapping Theorem.

P.T.O.



8. State and prove the Third Version of Cauchy's Theorem.
9. Prove the following : let G be a connected open set and let $f : G \rightarrow \mathbb{C}$ be an analytic function. Then the following conditions are equivalent.
- $f \equiv 0$;
 - there is a point a in G such that $f^n(a) = 0$ for each $n \geq 0$;
 - $\{z \in G : f(z) = 0\}$ has a limit point in G .

Unit – II

10. a) Show that for $a > 1$, Show that $\int_0^\pi \frac{d\theta}{a + \cos \theta} = \frac{\pi}{\sqrt{a^2 - 1}}$.
 b) State and prove the Residue theorem.
11. State and prove the Laurent Series Development.
12. Prove the following :
- If $|a| < 1$ then $\phi_a(z) = \frac{z-a}{1-\bar{a}z}$ is a one-one map of $D = \{z : |z| < 1\}$ on to itself ; the inverse of ϕ_a is ϕ_{-a} . Furthermore, ϕ_a maps ∂D on to ∂D , $\phi'_a(0) = 1 - |a|^2$ and $\phi'_a(a) = (1 - |a|^2)^{-1}$.
 - Let $f(z) = \frac{1}{z(z-1)(z-2)}$; give the Laurent series of $f(z)$ in each of the following annuli :
 - $\text{ann}(0 ; 0, 1)$,
 - $\text{ann}(0 ; 1, 2)$,
 - $\text{ann}(0 ; 2, \infty)$.

Unit – III

13. a) Prove the following : If G is open in $\mathbb{C}$ then there is a sequence $\{K_n\}$ of compact subsets of G such that $G = \bigcup_{n=1}^\infty K_n$. Moreover the sets K_n can be chosen to satisfy the following conditions :
- $K_n \subset \text{int } K_{n+1}$.
 - $K \subset G$ and K is compact implies $K \subset K_n$ for some n .
 - Every component of $G - K_n$ contains a component of $G - G$.
- b) State and prove Hurwitz's theorem.



14. a) With the usual notations, prove that $|1 - E_p(z)| \leq |z|^{p+1}$ for $|z| \leq 1$ and $p \geq 0$.
b) Discuss the convergence of the infinite product $\prod_{n=1}^{\infty} \frac{1}{n^p}$ for $p > 0$.
15. a) Show that $\prod (1 + z_n)$ converges absolutely iff $\prod (1 + |z_n|)$ converges.
b) Prove the following : If $\operatorname{Re} z_n > 0$ then the product $\prod z_n$ converges absolutely iff the series $\sum (z_n - 1)$ converges absolutely.
c) Prove the following : Let $\operatorname{Re} z_n > 0$ for all $n \geq 1$. Then $\prod_{n=1}^{\infty} z_n$ converges to a non zero number iff the series $\sum_{n=1}^{\infty} \log z_n$ converges.
-